

DISCRETE U-DUALITY GROUPS

Germar Schröder

*Max-Planck-Institut für Gravitationsphysik, Albert-Einstein-Institut,
Am Mühlenberg 1, D-14476 Golm, Germany*
germar@aei-potsdam.mpg.de

Abstract Generators for the discrete U-duality groups of toroidally compactified M-theory in $d \geq 4$ are presented and used to determine the $d = 3$ U-duality group. This contribution summarizes the results of [1].

1. INTRODUCTION AND SUMMARY

The study of nonperturbative duality symmetries in the last years has dramatically changed our understanding of string theory. U-duality, introduced in [2], is one of these symmetries. It was conjectured to be generated by the target-space duality of T^d (see [4] for a review) and the modular group of the torus including the eleventh M-theory direction [5]. This definition was adopted in the algebraic approach to U-duality reviewed in [3].

Using instead the original conjecture of [2] that takes the hidden symmetries of low energy supergravity as a starting point, generators of discrete U-duality in four dimensions may be determined directly. Groups in higher dimensions can be found by embeddings, and a method to determine the $d = 3$ U-duality is presented. Applied to a toy model corresponding to a truncation of M-theory, the method is seen to give a significantly different result than for the full theory.

2. U-DUALITY IN FOUR DIMENSIONS

Compactifying eleven dimensional supergravity, the low energy limit of M-theory, on T^7 yields 70 scalars arising from the $d = 11$ 3-form potential and the moduli of the torus. They may be joint in a field $\mathcal{V}^{(4)} \in E_{7(+7)}/SU(8)$ [7], a representation matrix in the fundamental **56** representation. Furthermore, 28 $U(1)$ gauge fields arise from the

$d = 11$ potential and metric. These fields and their magnetic duals can be arranged to 28 dimensional vectors $\mathcal{G}_{\bar{\mu}\bar{\nu}}$ and $\mathcal{H}_{\bar{\mu}\bar{\nu}}$, $\bar{\mu}, \bar{\nu} = 0 \dots 3$.

The equations of motion of the theory are invariant under classical $E_{7(+7)}$ duality and local $SU(8)$ transformations, acting as

$$\mathcal{F}_{\bar{\mu}\bar{\nu}} \rightarrow \Lambda^{-1} \mathcal{F}_{\bar{\mu}\bar{\nu}}, \quad \mathcal{V}^{(4)} \rightarrow h \mathcal{V}^{(4)} \Lambda, \quad \Lambda \in E_{7(+7)}, \quad h \in SU(8)$$

where $\mathcal{F}^t = (\mathcal{G}_{\bar{\mu}\bar{\nu}}, \mathcal{H}_{\bar{\mu}\bar{\nu}})^t$. Defining charges $\mathcal{Z} = \oint_{\Sigma} \mathcal{F}$, the DSZ quantization condition breaks $E_{7(+7)}$ to a discrete subgroup inducing integer shifts on the charge lattice. This group is the U-duality group and was proposed to extend to a nonperturbative quantum symmetry of M-theory.

To make U-duality transformations “manageable” in $d = 4$, the **56** representation needs to be addressed. This can be done by an embedding into $\mathfrak{e}_{8(+8)}$ using Freudenthal’s realization of exceptional Lie algebras [8]. The $\mathfrak{e}_{8(+8)}$ generators are given by E^i_j , $i, j = 1 \dots 9$, corresponding to $\mathfrak{sl}_9$, and E^{ijk} , E^*_{ijk} . Their commutators are given in [1]. Defining the basis

$$\mathcal{S}^t = \left(-E^*_{i\bar{j}9}, +E^1_{\bar{i}} \mid -E^{1\bar{i}\bar{j}}, -E^{\bar{i}}_9 \right), \quad \mathcal{X}^t = \left(x^{\bar{i}\bar{j}}, x^{\bar{i}9} \mid y_{\bar{i}\bar{j}}, y_{\bar{i}9} \right), \quad (1.1)$$

$\bar{i}, \bar{j} = 2 \dots 8$, the adjoint action of the $\mathfrak{e}_{7(+7)}$ subalgebra on $\mathcal{XS}$ exactly spans the **56** representation as defined in [7]. The basis (1.1) can therefore be used to study U-duality transformations on $\mathcal{F}$ and $\mathcal{V}^{(4)}$, as well as its subgroups T-duality and S-duality.

What are the generators of $E_{7(+7)}(\mathbb{Z})$? Using the fact that the **56** representation of $E_{7(+7)}$ is the unique minimal representation of $E_{7(+7)}$, it may be proven using the Birkhoff decomposition of Lie groups that the subgroup of $E_{7(+7)}$ inducing integer shifts on the lattice defined by $\mathcal{S}$ is generated by “fundamental unipotents”¹, that is, the action of the discrete subgroup is spanned by exponentiating the Chevalley generators for all positive and negative roots. From these, T and S generators may be built parallel to $SL(2, \mathbb{Z})$, the latter carrying a representation of the Weyl group modulo $\mathbb{Z}_2$. This together with the basis $\mathcal{S}$ yields contact to the algebraic approach to M-theory, and it may be shown that the two approaches are equivalent.

Since the notion of the above generators is representation independent, the discrete U-duality groups of higher dimensional theories follow directly from truncating the Dynkin diagram. Their representations are minimal and can be read off from $\mathcal{S}$.

¹See [9], where this group is defined in a more general context as homeomorphism of the group ring over $\mathbb{Z}$ to $\mathbb{Z}$.

2.1 U-DUALITY IN THREE DIMENSIONS

The $d = 3$ theory is known to have classical $E_{8(+8)}$ symmetry. As only scalars remain in the theory, the notion of electric charge seems ill defined and the meaning of a duality symmetry seems unclear. Therefore, in order to define U-duality in $d = 3$, a method proposed in [2] parallel to [6] may be extended to M-theory. By compactifying M-theory on the torus, we can choose eight different ways how to compactify first to four dimensions. This results in eight $E_{7(+7)}(\mathbb{Z})$'s acting *differently* on M-theory fields. By reducing the theory further to three dimensions, these groups are merged together to form the three dimensional duality group.

The reduction to $d = 3$ yields a scalar coset matrix of the form

$$\mathcal{V}^{(3)} = \mathcal{V}'^{(4)} \exp\left(\frac{1}{2}\phi \sum_{i=1}^8 h_i\right) \exp(\mathcal{Y} \cdot \mathcal{S}) \exp(f E_9^1). \quad (1.2)$$

$\mathcal{V}'^{(4)}$ is identical to $\mathcal{V}^{(4)}$, but now in the **248** adjoint representation of $E_{8(+8)}$. $\mathcal{Y}$ obeys $\partial_\mu \eta = \mathcal{G}_{\mu z}$, $\partial_\mu \bar{\eta} = \mathcal{H}_{\mu z}$, $\mathcal{Y} = (\eta, \bar{\eta})^t$, $\mu, \nu = 0 \dots 2$ and therefore carries the $d = 4$ charges, where z is the compact fourth direction. φ and f are the $d = 3$ dilaton and dualized KK field strength respectively, and h_i represents the $\mathfrak{sl}_9$ Cartan subalgebra.

A $\Lambda \in E_{7(+7)} \subset E_{8(+8)}$ transformation acts on $\mathcal{V}'^{(4)}$ and $\mathcal{Y}$ exactly as discussed in the last section. It therefore represents the $d = 4$ U-duality in $d = 3$. Completing a circle around vortex solutions in $d = 3$ may be seen to correspond to an $E_{7(+7)}(\mathbb{Z})$ action on the fields, parallel to [6].

The different compactifications yield 8 different coset matrices in $d = 3$. Using explicitly the connection to M-theory fields, it can be seen that they are related by

$$\mathcal{V}_{\#n}^{(3)} = (P_n S_n^1)^{-1} h_n \mathcal{V}_{\#1}^{(3)} P_n S_n^1$$

where h_n is a local transformation restoring upper triangular parameterization, S_n^1 corresponds to a Weyl reflection and P_n corresponds to a charge conjugation in $d = 4$. Denoting the $d = 4$ U-duality generators of the n th compactification by Λ , the total $d = 3$ U-duality is given by joining all generators

$$P_n S_n^1 \Lambda (P_n S_n^1)^{-1}$$

for all n . This can be seen to yield the whole $E_{8(+8)}(\mathbb{Z})$ defined by exponentiating all Chevalley generators. The intersection of two different $d = 4$ U-dualities is seen to be $E_{6(+6)}(\mathbb{Z})$ as expected. This determines U-duality in $d = 3$.

2.2 $G_{2(+2)}$ AS TOY MODEL

The described method to generate $d = 3$ U-duality can also be applied to five dimensional simple supergravity as toy model, which upon reduction to three dimensions exhibits a $G_{2(+2)}$ global symmetry [10]. It is known that this theory closely resembles $d = 11$ supergravity in many respects and corresponds to a direct truncation. The analogue of U-duality in $d = 4$ is $SL(2, \mathbb{Z})$, acting in the spin $3/2$ representation.

Here, the joint U-duality $U(\mathbb{Z})$ in $d = 3$ is strictly smaller than $G_{2(+2)}(\mathbb{Z})$ defined by exponentiating the Chevalley generators for all roots. All generators of $U(\mathbb{Z})$ correspond to short roots of $G_{2(+2)}(\mathbb{Z})$. That the groups do not agree is therefore connected to the fact that $G_{2(+2)}$ is not simply laced. Since no string compactification described by this no-moduli supergravity at low energies is known, one cannot determine which group is the “correct” U-duality group until such a description has been found.

Acknowledgments

I wish to thank Shun’ya Mizoguchi for a stimulating and fruitful collaboration.

References

- [1] S. Mizoguchi, G. Schröder, preprint hep-th/9909150.
- [2] C.M. Hull and P.K. Townsend, Nucl. Phys. **B438** (1995) 109.
- [3] N.A. Obers, B. Pioline, preprint hep-th/9809039.
- [4] A. Giveon, M. Porrati, E. Rabinovici, Phys. Rep. **244** (1994) 77.
- [5] J.H. Schwarz, Phys. Lett. **360B** (1995) 13; P. S. Aspinwall, Nucl. Phys. Proc. Supp. **46** (1996) 30; E. Witten, Nucl. Phys. **B443** (1995) 85.
- [6] A. Sen, Nucl. Phys. **B434** (1995) 179; A. Sen, Nucl. Phys. **B447** (1995) 62.
- [7] E. Cremmer and B. Julia, Nucl. Phys. **B159** (1979) 141; E. Cremmer, B. Julia, H. Lu, C.N. Pope, Nucl. Phys. **B523** (1998) 73.
- [8] H. Freudenthal, Proc. Kon. Ned. Akad. Wet. **A56** (Indagationes Math. **15**) (1953) 95-98 (French).
- [9] H. Matsumoto, Proceedings of Symposia in Pure Mathematics, American Mathematical Society, Vol. 9 (1966) 99; H. Matsumoto, Ann. scient. Éc. Norm. Sup., 4^e série **2** (1969) 1 (French).
- [10] S. Mizoguchi, N. Ohta, Phys. Lett. **441B** (1998) 123.